



GRANULAR FUNGICIDE V

Containing Chloroneb

- Provides quick knock down of *Pythium* blight.
- Contains chloroneb, an effective fungicide for controlling existing *Pythium* infestations.
- Prevents and controls grey snow mold.
- For use on all turfgrasses.
- May be applied by drop-type or rotary spreader.
- Choice of application rates, depending upon type and severity of infestation and degree of protection desired.
- Convenient, dry-applied, ready to use, free-flowing product with active ingredient impregnated on and into a granular carrier.

GENERAL INFORMATION

Bulk Density:	31.0 lb/cu. ft.
Net Weight:	14.86 kg (32.84 lb)
Coverage:	1,000 m ² (10,770 sq. ft.)

PRODUCT SPECIFICATIONS

Active ingredient: Chloroneb 6.25%

A dry, ready to use, free flowing product with active ingredient impregnated on and into a granular carrier.

GRANULAR FUNGICIDE V

Containing Chloroneb

REGISTRATION NUMBER 11466 PEST CONTROL PRODUCTS ACT

**FOR PROFESSIONAL USE ONLY.
KEEP OUT OF REACH OF CHILDREN.
READ THE LABEL BEFORE USING.**

DIRECTIONS FOR USE:

RECOMMENDED RATE: 14.86 kg/1,000 m²

DOUBLE RATE: 29 kg/1,000 m²

For use on turfgrass areas

Whenever used on greens and fairways apply to moist foliage in the morning or evening. Sprinkle lightly just before traffic resumes to avoid pickup of material on shoes, golf balls, or maintenance equipment.

To prevent Pythium blight – apply at the RECOMMENDED RATE to moist foliage when high temperatures and humidity favour development. Repeat at 5-7 day intervals as long as these weather conditions persist.

To control Pythium blight – apply at the RECOMMENDED RATE to moist foliage as soon as disease activity is observed; repeat at 5-7 day intervals during high temperature and humidity conditions.

In case of severe disease attack apply at the DOUBLE RATE to moist foliage and continue RECOMMENDED RATE applications at 5-7 day intervals during high temperature and humidity conditions.

On new seedlings, apply at the RECOMMENDED RATE after sowing seed; at first signs of disease apply to moist foliage of newly germinated grass, and repeat at 7-10 day intervals if evidence of disease persists.

When treating to control Pythium blight do not fertilize or water heavily until disease activity is arrested; prolonged contact with foliage enhances effectiveness.

Grey snow mold – apply at the RECOMMENDED RATE to dry or moist foliage prior to first heavy snow (early winter). Repeat in midwinter if turf is free of snow. A final treatment in late winter as the snow melts, extends protection.

In areas where snow coverage is present the entire winter, apply at the DOUBLE RATE prior to the first snow and at the RECOMMENDED RATE when snow melts in the spring.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. May irritate eyes, nose, throat and skin. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not graze or feed clippings from treated areas to livestock. Do not contaminate water when disposing of equipment washwaters.

FIRST AID:

IF ON SKIN: Wash with plenty of water.

IF IN EYES: Flush with water for 5-10 minutes and get medical attention or contact poison control centre IMMEDIATELY. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

DISPOSAL:

1. Thoroughly empty the contents of the container into the application device.
2. Make the empty container unsuitable for further use.
3. Dispose of the container in accordance with provincial requirements.
4. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

SUGGESTED SPREADER SETTINGS

To provide proper distribution, calibrate spreader before application
14.86 kg treats 1,022 m² at RECOMMENDED RATE (Use 14.86 kg/1,000 m²)
14.86 kg treats 511 m² at DOUBLE RATE (Use 29 kg/1,000 m²)

SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS	
			RECOMMENDED RATE	DOUBLE RATE
ProTurf® Drop	5 km/hr	Overlap Wheels	6	8
ProTurf® Rotaries R-7, R7X, R-8, R8-A (cone 8)	5 km/hr	2.4 m	G	K
	5 km/hr	2.7 m	M	Q
Accupro 2000 (cone 9)	5 km/hr	3.0 m	K	O 1/2
SR 2000 (cone 9)	5 km/hr	3.0 m	K	O 1/2
Lely-WTR and WRF Models	7 km/hr	4.6 m	4 1/2 III	6 III
Lely-HR (PTO Model)	450 rpm @ 7 km/hr	4.9 m	4 1/2 III	6 III



Always refer to the actual product label for complete directions and precautions.

ProTurf® is a registered trademark, used under licence from The Scotts Company.
Nu-Gro IP Inc., 10 Craig Street, Brantford, Ontario N3R 7J1

APRIL, 2003