

# QUALI-PRO



Department of Agriculture  
STATE OF HAWAII

LICENSED

**8275.69**

PERIOD 2011-2013 LIC. NO.

# Tebuconazole 3.6F

## *Foliar Fungicide*

### TURF AND ORNAMENTAL USES\*

\*Not for use in California

### ACTIVE INGREDIENT:

% BY WT.

Tebuconazole: alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol . . . . . 38.7%

OTHER INGREDIENTS: . . . . . 61.3%

TOTAL: . . . . . 100.0%

Contains 3.6 pounds tebuconazole per gallon

EPA Reg. No. 66222-117

EPA Est. No. 37429-GA-001<sup>BT</sup>; 37429-GA-002<sup>BO</sup>

Letter(s) in lot number correspond(s) to superscript in EPA Est. No.

## KEEP OUT OF REACH OF CHILDREN

# CAUTION

For additional precautionary, handling, and use statements,  
see inside of this booklet.

Manufactured for: **Makhteshim Agan of North America, Inc.**  
4515 Falls of Neuse Road, Suite 300 • Raleigh, NC 27609

13931

EPA 032411/Rev A

## Net Contents: 1 Gallon

### FIRST AID

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For medical emergencies, call Prosar 24 hours a day at 1-877-250-9291.

**NOTE TO PHYSICIAN:** No specific antidote. Treat symptomatically.

**Symptoms of Poisoning:** The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride, or Viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washing exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### USER SAFETY RECOMMENDATIONS

**Users should:**

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish, and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

**Ground Water Advisory:** Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**Surface Water Label Advisory:** This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours for all crops.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride, or Viton
- Shoes plus socks

#### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**Entry Restrictions for Non-WPS Uses:** Do not enter or allow others to enter the treated area until sprays have dried.

**Spray Volume:** For turf, apply Quali-Pro Tebuconazole 3.6F in 66-132 gallons of water per acre by ground sprayer. For ornamentals other than leatherleaf fern, use 50-300 gallons of finished spray per acre depending upon equipment, plant species, and plant growth stage at the time of application. For leatherleaf fern, apply Quali-Pro Tebuconazole 3.6F in a minimum of 5 gallons of finished spray per acre using ground equipment or chemigation. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.

**Chemigation:** Apply Quali-Pro Tebuconazole 3.6F through irrigation equipment only to leatherleaf fern in Florida to suppress anthracnose. Apply this product only through center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move; or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. Contact State Extension Service specialists, equipment manufacturers, or other experts if you have questions regarding calibration. Do not connect irrigation systems (including

greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments if the need arises.

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection.

The system must contain a functional check valve, vacuum-relief valve, and low-pressure drain, appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. Pesticide may be applied continuously for the duration of the water application.

**Mixing:** Add labeled amount of Quali-Pro Tebuconazole 3.6F into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the Quali-Pro Tebuconazole 3.6F should be thoroughly dispersed prior to the addition of other materials.

Do not tank mix with products containing a prohibition against tank mixing. Follow the most restrictive labeling requirements of any tank mix product.

**Compatibility:** To determine the compatibility of Quali-Pro Tebuconazole 3.6F with other products, use the following procedure: Pour the labeled proportions of the products into a suitable container of water, mix thoroughly, and allow to stand at least five (5) minutes. If the combination remains mixed or can be remixed readily, the mixture is considered physically compatible. For further information, contact your local Makhteshim Agan representative.

#### **Resistance Management Statement**

Quali-Pro Tebuconazole 3.6F is a Group 3 fungicide which exhibits no known cross-resistance to other fungicide groups. However, fungal pathogens are known to develop resistance to products with the same mode of action when used repeatedly. Any fungal population may contain or develop individuals that are resistant to Quali-Pro Tebuconazole 3.6F and other Group 3 fungicides. If Group 3 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted diseases, the resistance isolates may eventually dominate the fungal population. Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include rotation and/or tank mixing with products having different modes of action or limiting the total number of applications per season. Contact your local extension specialist, certified crop advisor, and/or manufacturer for fungicide resistance management and/or integrated disease management recommendations for specific crops and resistant disease populations. Makhteshim Agan of North America, Inc. encourages responsible management to ensure effective long-term control of the fungal disease on this label.

### **TURF AND ORNAMENTAL USES**

#### **DISEASE CONTROL IN GOLF COURSE TURF**

#### **PRODUCT INFORMATION**

For use on all golf turf applications of cool-season and warm-season grasses (such as Bentgrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, and Zoysia) or their mixtures. Quali-Pro Tebuconazole 3.6F is not phytotoxic to any of the above-mentioned grasses when used in accordance with the label. **Note:** Bermudagrass can be sensitive to Quali-Pro Tebuconazole 3.6F under certain conditions. Do not apply consecutive applications during or just after dormancy break. Avoid applications when temperatures are expected to exceed 85° F.

Quali-Pro Tebuconazole 3.6F can be used for the prevention and control of the diseases mentioned in table below. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Preventative treatments can be applied using 28-day intervals as indicated. When treating golf greens, always treat aprons and approaches. Spray uniformly over the area to be treated with properly calibrated equipment. Apply the specified amount of Quali-Pro Tebuconazole 3.6F Fungicide in sufficient water for thorough coverage. A volume of 66 – 132 gallons per acre (1.5 – 3.0 gallons per 1000 sq ft) is recommended. Apply using properly calibrated low-volume, hand-held, mechanical, or motorized ground broadcast equipment. Application to small areas may be made with low-pressure handwand or backpack equipment.

Depending on the disease, Quali-Pro Tebuconazole 3.6F should be watered into the crown and active root zone for best results. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. For best results, use spray mixture the same day it is prepared.

#### USE RESTRICTIONS AND PRECAUTIONS

- For use on golf course turf only.
- Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle, and high school), campgrounds, churches, and theme parks.
- Not for homeowner use.
- Not for use on turf being grown for sale or commercial use as sod.
- Do not use clippings for animal feed.
- Do not exceed 3.6 oz of Quali-Pro Tebuconazole 3.6F Fungicide per 1000 sq ft per year.
- Do not apply more than 6 applications per year.

DISEASE	RATE OF QUALI-PRO TEBUCONAZOLE 3.6F FL OZ/1000 SQ FT	REMARKS
Dollar Spot ( <i>Sclerotinia homoeocarpa</i> ) Copper Spot ( <i>Gloeocercospora sorghi</i> ) Powdery Mildew ( <i>Erysiphe graminis</i> ) Corticium Red Thread ( <i>Laetisaria fuciformis</i> ) Rusts ( <i>Puccinia</i> spp.)	0.6	For prevention, begin applications when conditions are favorable for disease development. Do not make two consecutive applications of Quali-Pro Tebuconazole 3.6F. Alternate with another fungicide with a different mode of action. A second application may be made after 28 days.
Brown Patch/Rhizoctonia Blight, Large Patch ( <i>Rhizoctonia solani</i> ) Brown Ring Patch ( <i>R. circinata</i> )	0.6	
Anthraxnose-Basal and Foliar ( <i>Colletotrichum cereale</i> ) Red Thread ( <i>Laetisaria fuciformis</i> ) Pink Patch ( <i>Limonomyces rosipellis</i> )	0.6	

See COMMENTS at end of table.

DISEASE	RATE OF QUALI-PRO TEBUCONAZOLE 3.6F FL OZ/1000 SQ FT	REMARKS
<b>Bermudagrass decline</b> <i>(Gaeumannomyces graminis var. graminis)</i>	0.6	<p>Immediately after fungicide is applied, the area should be irrigated with sufficient water to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone.</p> <p>For prevention, begin applications two or four weeks prior to the historical appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Apply subsequent applications at 28-day intervals.</p>
<b>Take All Patch</b> <i>(Gaeumannomyces graminis)</i>	0.6	<p>For prevention, apply in the fall when soil temperature reaches 55-65° F and again in the spring under similar soil temperature conditions. Applications in both fall and spring may be necessary.</p> <p>Immediately after fungicide is applied, the area should be irrigated with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.</p>
<b>Gray Leaf Spot</b> <i>(Pyricularia grisea)</i>	0.6	<p>Apply when conditions are favorable for disease development at 28-day intervals. If using 0.6 oz/1000 sq ft, or under conditions favoring moderate to heavy disease pressure, Quali-Pro Tebuconazole 3.6F can be tank mixed with a registered contact fungicide at label rate.</p>
<b>Stripe Smut</b> ( <i>Ustilago striiformis</i> )	0.6	<p>Make a single application to historical disease areas in spring as grass growth begins.</p>

See **COMMENTS** at end of table.



DISEASE	RATE OF QUALI-PRO TEBUCONAZOLE 3.6F FL OZ/1000 SQ FT	REMARKS
Spring Dead Spot ( <i>Leptosphaeria korrea</i> , <i>L. narmari</i> , <i>Ophiosphaerella her- potricha</i> , <i>Gaeumannomyces graminis</i> ) Necrotic Ring Spot ( <i>Leptosphaeria korrea</i> )	0.6	For prevention, apply in fall when soil temperature reaches 65° F and again in spring under similar soil temperature conditions or after dormancy break. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Fusarium Patch ( <i>Fusarium roseum</i> )	0.6	Apply first application in mid-June or 28 days prior to time this blight normally becomes evident. Make applications at no less than 28-day intervals.
Summer Patch ( <i>Magnaporthe poae</i> )	0.6	Apply beginning in the spring. Do not make two consecutive applications of Quali-Pro Tebuconazole 3.6F. Alternate with another fungicide with a different mode of action. Second and third applications may be made at 28-day intervals. See local university recommendations for suggested timing. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
Zoysia Patch, Large Patch of Zoysia ( <i>Rhizoctonia solani</i> )	0.6	Make first application in early fall (mid-September to mid-October) prior to development of disease symptoms. A second application in early spring may be necessary in areas where disease pressure is known to be heavy.

See COMMENTS at end of table.

DISEASE	RATE OF QUALI-PRO TEBUCONAZOLE 3.6F FL OZ/1000 SQ FT	REMARKS
<p><b>Gray Snow Mold/Typhula Blight</b> (<i>Typhula incarnate</i>)</p> <p><b>Pink Snow Mold/Microdochium Patch</b> (<i>Microdochium nivale</i>)</p>	0.6	Apply in the fall, before anticipated turf dormancy and before first snow cover. If turf breaks dormancy during winter months, a second application may be made. Do not apply over snow cover, or when turf is dormant. It is recommended that Quali-Pro Tebuconazole 3.6F be tank mixed with other registered snow mold products for best season-long results.
<p><b>COMMENTS:</b> Apply the specified amount of Quali-Pro Tebuconazole 3.6F in 1.5 to 3.0 gallons of water per 1000 sq ft. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. Do not use clippings for animal feed. Do not exceed 3.6 fl oz of Quali-Pro Tebuconazole 3.6F per 1000 sq ft per year. Do not exceed 6 applications per year.</p>		

**DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS, AND COMMERCIAL AND RESIDENTIAL LANDSCAPES**

**PRODUCT INFORMATION:** For use on ornamental plants only; not for woodlands or forest management. Not for homeowner use.

Quali-Pro Tebuconazole 3.6F can be used in a preventative and curative disease control program for the listed plant types and diseases in the table below. Optimum disease management is obtained when Quali-Pro Tebuconazole 3.6 F is used in conjunction with sound disease management practices.

Apply material with properly calibrated hand-held, mechanical, or motorized spray equipment. Begin applications when disease first appears and repeat at 14- to 21-day intervals during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand-held, mechanical, or motorized applications, mix as directed below and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Choose a finished spray volume appropriate for the size of the plants and amount of foliage which will provide thorough coverage throughout the canopy. Allow sprays to dry before overhead irrigation is applied.

**USE RESTRICTIONS AND PRECAUTIONS**

Apply Quali-Pro Tebuconazole 3.6F at rates of 4-10 fl oz per acre in 100 gallons of water. Spray volume may range from 50 up to 300 gallons of finished spray per acre depending upon equip-

- ment, plant species, and plant growth stage at time of application.
- Do not apply more than 30 fl oz per acre in a single application.
- Do not apply more than 0.94 gallons (120 fl oz) of Quali-Pro Tebuconazole 3.6F (equal to 3.38 lbs of tebuconazole) per acre per year.
- Do not make more than 4 applications per year at highest rate.
- Do not apply to bearing fruit trees or vegetables.

**NOTE:** The **DIRECTIONS FOR USE** section of this product label reflects the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on African Violets, Begonias, Boston Fern, and Geraniums.

#### ORNAMENTALS DISEASE CONTROL

PLANTS	DISEASE	APPLICATION	
		TO PREVENT DISEASE	TO TREAT DISEASE
<b>Roses</b>	Black Spot Powdery Mildew Rust	Apply every 14-21 days during the growing season, starting when leaves first appear.	Apply every 14 days for a total of 3 applications beginning at the first sign of disease.
<b>Flowers</b>	Leaf Spot Powdery Mildew Rust Southern Blight	Apply at least 3 times per year, 14-21 days apart, beginning with spring bud break.	
<b>Crabapples (Ornamental), Dogwoods, and other Landscape (Ornamental) Trees</b>	Anthracnose Leaf Spot Powdery Mildew Rust Scab	Rotation or tank mixing with barrier protectant fungicides is recommended for resistance management.	
<b>Azaleas, Camellias, Rhododendrons, and other Landscape (Ornamental) Shrubs</b>	Anthracnose Black Spot Leaf Spot Petals Blight Powdery Mildew	Petals Blight: Apply 2-3 times per week into the flowers as they open and develop color.	
<b>Ground Covers and Vines</b>	Rust Southern Blight		

For small plantings, add 1 teaspoon of Quali-Pro Tebuconazole 3.6F to 2.5 gallons of water.

### Pump-Style Sprayers

1. Add the appropriate amounts of concentrate and water to the sprayer tank.
2. Close the sprayer, shake well, and pressurize.
3. Adjust nozzle to a coarse spray pattern and apply.
4. Occasionally repressurize the sprayer, if needed, to maintain a good spray pattern.

PLANT	DISEASE	RATE OF QUALI-PRO TEBUCONAZOLE 3.6 F
LEATHERLEAF FERN (FLORIDA ONLY)	Anthracnose (suppression)	5 to 10 fl oz per acre
	<b>Notes:</b> Make the first application before anthracnose symptoms are present and continue at 12- to 14-day intervals. <b>USE RESTRICTIONS:</b> A maximum of 5 pints of Quali-Pro Tebuconazole 3.6F may be applied per acre per year.	

**Comments:** Apply Quali-Pro Tebuconazole 3.6F in a minimum of 5 gallons of spray solution per acre using ground equipment or chemigation.

#### USE LIMITATIONS:

Quali-Pro Tebuconazole 3.6F can cause phytotoxicity to Leatherleaf Fern under certain environmental conditions. Applications in temperatures less than 70°F can cause phytotoxicity in the form of leaf burning and/or yellowing. Application followed by temperatures falling below 55°F can cause similar symptoms.

Before using this product on Leatherleaf Fern, read the **LIMITATION OF WARRANTY AND LIABILITY** section in its entirety.

### OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES, OR NATURAL PONDS, AND ESTUARIES.

- Apply only during alternate years in fields adjacent to aquatic areas listed above.
- Do not apply by ground or air within 100 feet of aquatic areas listed above.
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

**Spray Drift Management:** For aerial applications, mount the spray boom on the aircraft so as to minimize drift caused by wing tip vortices. Use the minimum practical boom length and do not exceed 75% of the wing span or rotor diameter. Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the airstream as much as possible, and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Release the spray at the lowest possible height consistent with good pest control and flight safety. Avoid applications more than 10 feet above the crop canopy. Make aerial or ground appli-

cautions when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area. Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature. Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking, invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to **PRECAUTIONARY STATEMENTS** on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing-type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

### CONTAINER HANDLING:

**Nonrefillable Container (five gallons or less):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinse into application equipment or a mix tank or store rinseate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

## LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES and LIMITATIONS OF LIABILITY.**

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this prod-

uct. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Makhteshim Agan of North America, Inc.'s election, the replacement of product.

To the extent consistent with applicable law, MANA accepts no responsibility and shall not be liable for phytotoxicity or side effects of Quali-Pro Tebuconazole 3.6F used on Leatherleaf ferns under any conditions.

Quali-Pro is a registered trademark of Makhteshim Agan of North America, Inc.

# Quali-Pro Tebuconazole 3.6F

## Foliar Fungicide

Contains tebuconazole, the active ingredient used in Folicur 3.6F.  
Quali-Pro Tebuconazole 3.6F is not manufactured or distributed by Bayer CropScience.

<b>ACTIVE INGREDIENT:</b>	<b>% BY WT.</b>
Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol .....	38.7%
<b>INERT INGREDIENTS:</b> .....	<u>61.3%</u>
	<b>Total: 100.0%</b>



**LICENSED**

PERIOD 2011-2013 LIC. NO.

**8275.69**

Contains 3.6 pounds tebuconazole per gallon  
**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

Manufactured for:  
Makhteshim Agan of North America, Inc.  
4515 Falls of Neuse Rd., Suite 300  
Raleigh, NC 27609

EPA Reg. No. 66222-117

EPA Est. No. 37429-GA-001

**NET CONTENTS: 2.5 GALLONS**

### FIRST AID

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For medical emergencies, call Prozar 24 hours a day at 1-877-250-9291. <b>NOTE TO PHYSICIAN:</b> No specific antidote. Treat symptomatically. <b>Symptoms of Poisoning:</b> The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.</p>	

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride, or Viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

**Users should:**

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish, and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

**Ground Water Advisory:** Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**Surface Water Label Advisory:** This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours for all crops except sweet corn and lychee. For sweet corn, the REI is 19 days. For lychee, the REI is 2 days.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride, or Viton
- Shoes plus socks

**Spray Volume:** Quali-Pro Tebuconazole 3.6F may be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.

**Chemigation:** Do not apply this product through any type of irrigation system.

**Mixing:** Add labeled amount of Quali-Pro Tebuconazole 3.6F into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the Quali-Pro Tebuconazole 3.6F should be thoroughly dispersed prior to the addition of other materials.

Do not tank mix with products containing a prohibition against tank mixing. Follow the most restrictive labeling requirements of any tank mix product.

**Compatibility:** To determine the compatibility of Quali-Pro Tebuconazole 3.6F with other products, the following procedure should be followed: Pour the labeled proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least five (5) minutes. If the combination remains mixed or can be remixed readily, the mixture is considered physically compatible. For further information contact your local Makhteshim Agan representative.

### APPLICATION INSTRUCTIONS

CROP	DISEASE	RATE OF QUALI-PRO TEBUCONAZOLE 3.6 F
ASPARAGUS	Rusts ( <i>Puccinia spp.</i> )	4 to 6 fl oz per acre
	<p><b>Notes:</b> Apply Quali-Pro Tebuconazole 3.6F as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl oz of Quali-Pro Tebuconazole 3.6F per acre (0.11 lb ai – 0.17 lb ai per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher rate. Repeat applications on a 14-day interval as necessary to maintain control of rust. Do not apply to harvestable spears. Do not apply within 100 days of harvest in California and 180 days in all other states. Do not make more than three foliar applications per season (18 fl oz/acre or 0.51 lb ai/acre).</p>	
<p><b>General Comments:</b> Applications may be made using ground or aerial application equipment. A 50 foot spray drift buffer zone is required for all aerial applications. For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Quali-Pro Tebuconazole 3.6F. Quali-Pro Tebuconazole 3.6F is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating Quali-Pro Tebuconazole 3.6F with other DMI fungicides may lead to resistance. Restricted-entry interval (REI) = 12 hours.</p>		



CROP	DISEASE	RATE OF QUALI-PRO TEBUCONAZOLE 3.6 F
<b>BARLEY</b>	Rusts ( <i>Puccinia</i> spp.) Head blight ( <i>Fusarium</i> spp.) – Suppression	4 fl oz per acre
	<p><b>Notes:</b> Apply Quali-Pro Tebuconazole 3.6F in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. A maximum of 4 fl. oz. of Quali-Pro Tebuconazole 3.6F may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of Quali-Pro Tebuconazole 3.6F. Barley fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.</p> <p><b>Application timing directions:</b>  Rusts: Apply Quali-Pro Tebuconazole 3.6F at the earliest sign of rust pustules on foliage.  Fusarium head blight: Optimal timing of Quali-Pro Tebuconazole 3.6F for Fusarium head blight suppression is when main stem heads have fully emerged (Feekees 10.5) on 50% of the plants.</p>	
<p><b>General Comments:</b> For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Quali-Pro Tebuconazole 3.6F. Quali-Pro Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Quali-Pro Tebuconazole 3.6F will be resistant to weathering. Quali-Pro Tebuconazole 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).</p>		
<b>BEANS</b> (fresh & dry except succulent shelled)	Rust ( <i>Uromyces appendiculatus</i> )	4 to 6 fl oz per acre
	<p><b>Notes:</b> Apply Quali-Pro Tebuconazole 3.6F in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control. Beans, fresh: Quali-Pro Tebuconazole 3.6F may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Quali-Pro Tebuconazole 3.6F per acre per crop season. Beans, dry: Quali-Pro Tebuconazole 3.6F may be applied up to 14 days before harvest. Do not apply more than 12 fl. oz. of Quali-Pro Tebuconazole 3.6F per acre per crop season.</p>	
<p><b>General Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Quali-Pro Tebuconazole 3.6F. Quali-Pro Tebuconazole 3.6F must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Quali-Pro Tebuconazole 3.6F will be resistant to weathering. Quali-Pro Tebuconazole 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).  Restricted-entry interval (REI) = 12 hours.</p>		
<b>CORN</b> (sweet corn, field corn, field corn grown for seed, and popcorn)	Rust ( <i>Puccinia</i> spp.)  Northern leaf blight ( <i>Helminthosporium turcicum</i> )  Southern leaf blight ( <i>Helminthosporium maydis</i> )  Northern leaf spot ( <i>Helminthosporium carbonum</i> )  Gray leaf spot ( <i>Cercospora zea-maydis</i> )	4 to 6 fl oz per acre
	<p><b>Notes:</b> Apply Quali-Pro Tebuconazole 3.6F in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. A maximum of 24 fl. oz. (1.5 pint) of Quali-Pro Tebuconazole 3.6F may be applied per acre per crop season. Sweet corn: Quali-Pro Tebuconazole 3.6F may be applied up to 7 days before the harvest of ears or forage, and 49 days before the harvest of fodder. Field, seed, or popcorn: Quali-Pro Tebuconazole 3.6F may be applied up to 21 days before the harvest of forage, and 36 days before the harvest of grain or fodder.</p>	
<p><b>General Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Quali-Pro Tebuconazole 3.6F. Quali-Pro Tebuconazole 3.6F must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Quali-Pro Tebuconazole 3.6F will be resistant to weathering. Quali-Pro Tebuconazole 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).  Restricted-entry interval (REI) for sweet corn = 19 days.  Restricted-entry interval (REI) for all corn except sweet corn = 12 hours.</p>		
<b>COTTON</b>	Southwestern cotton rust ( <i>Puccinia cacabata</i> )	6 to 8 fl oz per acre
	<p><b>Notes:</b> Apply Quali-Pro Tebuconazole 3.6F in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. Quali-Pro Tebuconazole 3.6F may be applied up to 30 days before harvest. Do not apply more than 24 fl. oz. of Quali-Pro Tebuconazole 3.6F per acre per crop season.</p>	
<p><b>General Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Quali-Pro Tebuconazole 3.6F. Quali-Pro Tebuconazole 3.6F must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Quali-Pro Tebuconazole 3.6F will be resistant to weathering. Quali-Pro Tebuconazole 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).  Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF QUALI-PRO TEBUCONAZOLE 3.6 F
<b>CUCURBIT VEGETABLES GROUP</b> Chayote Chinese waxgourd Citron melon Cucumber Gherkin Edible gourd, (includes hyotan, cucuzza, hechima and Chinese okra) Momordica spp. (includes balsam apple, balsam pear, bitter melon and Chinese cucumber) Muskmelon (includes cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon) Pumpkin Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow and zucchini) Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash and spaghetti squash) Watermelon	Powdery mildew ( <i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i> ) ( <i>Erysiphe cichoracearum</i> )	4 to 6 fl oz per acre
	Gummy stem blight - suppression ( <i>Didymella bryonae</i> ) (watermelon, squash, pumpkin, and melons only)	8 fl oz per acre
<p><b>Notes:</b> Apply the specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals. Quali-Pro Tebuconazole 3.6F may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Quali-Pro Tebuconazole 3.6F per acre per crop season.</p>		
<p><b>General Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Quali-Pro Tebuconazole 3.6F. Quali-Pro Tebuconazole 3.6F must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Quali-Pro Tebuconazole 3.6F will be resistant to weathering. Quali-Pro Tebuconazole 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).            Restricted-entry interval (REI) = 12 hours.</p>		
<b>DRY BULB ONION</b> <b>GARLIC</b> <b>GREAT-HEADED (ELEPHANT) GARLIC</b> <b>WELCH ONION</b> <b>SHALLOT</b>	White rot ( <i>Sclerotium cepivorum</i> )	White rot: 20.5 fl oz per acre applied in a 4 to 6 inch band over/into each furrow. May be applied by chemigation to control white rot.
	Rust ( <i>Puccinia allii</i> , <i>Puccinia porri</i> ) Purple blotch ( <i>Alternaria porii</i> )	4 to 6 fl oz per acre
	<p>White rot: For the control of white rot, make one application in the furrow at the time of planting. The in-furrow application should be made at the rate of 20.5 fl. oz Quali-Pro Tebuconazole 3.6 F per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 to 6 fl oz/acre.</p> <p>Rust: For the control of rust make foliar applications at the rate of 4 to 6 fl. oz Quali-Pro Tebuconazole 3.6 F per acre per application. Repeat at an interval of 10 to 14 days.</p> <p>Apply Quali-Pro Tebuconazole 3.6F F in a protective spray schedule or when weather conditions are favorable for rust development.</p> <p><b>Notes:</b> Do not apply more than 32.5 fl. oz. Quali-Pro Tebuconazole 3.6 F per acre per season if an in-furrow treatment is made. If Quali-Pro Tebuconazole 3.6 F is not applied as an in-furrow treatment then do not apply more than 12 fl oz Quali-Pro Tebuconazole 3.6 F per acre per season as a foliar spray. Do not apply within 7 days of harvest (PHI = 7 days).</p>	
<p><b>General Comments:</b> For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant may be tank-mixed with Quali-Pro Tebuconazole 3.6F. Quali-Pro Tebuconazole 3.6F must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Quali-Pro Tebuconazole 3.6F will be resistant to weathering. Quali-Pro Tebuconazole 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).            Restricted-entry interval (REI) = 12 hours.</p>		
<b>GREEN ONION, LEEK, SPRING ONION, SCALLION, JAPANESE BUNCHING ONION, GREEN SHALLOTS AND GREEN ESCHALOTS</b>	White rot caused by <i>Sclerotium cepivorum</i> suppression only Rust ( <i>Puccinia allii</i> , <i>Puccinia porri</i> ) Purple blotch ( <i>Alternaria porii</i> )	4 to 6 fl oz per acre
	<p>For the control of diseases make foliar applications using an interval of 10 to 14 days. Apply Quali-Pro Tebuconazole 3.6F in a protective spray schedule or when weather conditions are favorable for rust development.</p> <p><b>Notes:</b> Do not apply more than 24 fl oz of Quali-Pro Tebuconazole 3.6F per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).</p>	
<p><b>General Comments:</b> For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant may be tank-mixed with Quali-Pro Tebuconazole 3.6F. Quali-Pro Tebuconazole 3.6F must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Quali-Pro Tebuconazole 3.6F will be resistant to weathering. Quali-Pro Tebuconazole 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).            Restricted entry interval (REI) is 12 hours.</p>		

CROP	DISEASE	RATE OF QUALI-PRO TEBUCONAZOLE 3.6 F
GARDEN BEET roots and tops (leaves)	Cercospora leaf spot ( <i>Cercospora beticola</i> )	3 to 7.2 fl oz per acre
	<b>Notes:</b> Make applications on 14 day intervals. Do not apply more than 28.8 fl. oz. Quali-Pro Tebuconazole 3.6F per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).	
<b>General Comments:</b> For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant may be tank-mixed with Quali-Pro Tebuconazole 3.6F. Quali-Pro Tebuconazole 3.6F must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Quali-Pro Tebuconazole 3.6F will be resistant to weathering. Quali-Pro Tebuconazole 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		
GRASSES GROWN FOR SEED	Rusts ( <i>Puccinia spp.</i> )	4 to 8 fl oz per acre
	Apply the specified rate of Quali-Pro Tebuconazole 3.6F as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.	
	Powdery mildew	4 to 8 fl oz per acre
	Apply specified rate of Quali-Pro Tebuconazole 3.6F when powdery mildew first appears on the leaves. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.	
<b>General Comments:</b> Apply the specified rate in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control. For optimum benefit, the lowest labeled rate of a spray surfactant should be tank mixed with Quali-Pro Tebuconazole 3.6F. A maximum of 16 fluid ounces (1 pint) may be applied per acre per crop season. Quali-Pro Tebuconazole 3.6F may be applied up to 4 days before harvest. Chaff, screenings and straw from treated areas may be used for feed purposes; however, do not use forage, cut green crop, or use seed for feed purposes. Regrowth may be grazed starting 17 days after last application.		
HOPS	Powdery mildew ( <i>Sphaerotheca humuli</i> / <i>Sphaerotheca macularis</i> )	4 to 8 fl oz per acre
	<b>Notes:</b> Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 10- to 14-day intervals. Quali-Pro Tebuconazole 3.6F may be applied up to 14 days before harvest. Do not apply more than 32 fl. oz. of Quali-Pro Tebuconazole 3.6F per acre per crop season. Increase the spray volume and the application rate as vine growth increases during the season.	
<b>General Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Quali-Pro Tebuconazole 3.6F. Quali-Pro Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Quali-Pro Tebuconazole 3.6F will be resistant to weathering. Quali-Pro Tebuconazole 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		
LEAFY BRASSICA GREENS (Broccoli raab, Chinese cabbage (bok choy), collards, kale, mizuma, mustard greens, mustard spinach, rape greens, turnip greens)	Cercospora leaf spot ( <i>Cercospora brassicicola</i> ) Powdery mildew ( <i>Erysiphe cruciferarum</i> ) Alternaria leaf spot ( <i>Alternaria brassicicola</i> )	3 to 4 fl oz
	<b>Notes:</b> Do not apply more than 16 fl oz Quali-Pro Tebuconazole 3.6F per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).	
<b>General Comments:</b> For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant may be tank-mixed with Quali-Pro Tebuconazole 3.6F. Quali-Pro Tebuconazole 3.6F must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Quali-Pro Tebuconazole 3.6F will be resistant to weathering. Quali-Pro Tebuconazole 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3). <b>Restriction:</b> Application to turnip greens is limited to East of the Rockies. Restricted Entry Interval (REI) is 12 hours.		
LYCHEE	Anthrachnose ( <i>Colletotrichum gloeosporioides</i> )	4 to 6 fl. oz. per acre
	<b>Notes:</b> Begin first application of Quali-Pro Tebuconazole 3.6F as panicle emerges. Spray up to 6 fl. oz. per acre every 10 days thereafter for a total of 8 sprays. Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only. Do not apply more than 48 fl. oz. of Quali-Pro Tebuconazole 3.6F per acre per season. Quali-Pro Tebuconazole 3.6F can be applied up to and including the day of harvest (PHI = 0 days).	
<b>General Comments:</b> For optimum disease control, the lowest labeled rate of a non-ionic spray surfactant should be tank-mixed with Quali-Pro Tebuconazole 3.6F. Quali-Pro Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Quali-Pro Tebuconazole 3.6F will be resistant to weathering. Quali-Pro Tebuconazole 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 2 days.		
MANGO (POST-HARVEST)	Anthrachnose ( <i>Colletotrichum gloeosporioides</i> )	7.5 fl. oz. per 100 gal. water / 50,000 lbs. fruit
	<b>Notes:</b> Quali-Pro Tebuconazole 3.6F can be applied as a post-harvest treatment by mixing at the specified rate. Application to the fruit is made by spraying the dilute Quali-Pro Tebuconazole 3.6F solution onto brushes in a commercial packing line.	

CROP	DISEASE	RATE OF QUALI-PRO TEBUCONAZOLE 3.6 F
OKRA	Cercospora leaf spot ( <i>Cercospora spp.</i> )	4 to 6 fl. oz per acre
	<b>Notes:</b> Apply specific dosage of Quali-Pro Tebuconazole 3.6F in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Applications may be made no closer than 3 days before harvest. Do not apply more than 24 fl. oz. of Quali-Pro Tebuconazole 3.6F per acre per season.	
<b>General Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Quali-Pro Tebuconazole 3.6F. Quali-Pro Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Quali-Pro Tebuconazole 3.6F will be resistant to weathering. Quali-Pro Tebuconazole 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		
PEANUTS	<b>SOILBORNE:</b> Sclerotium stem and pod rot (white mold, southern blight, southern stem rot) Rhizoctonia limb rot Rhizoctonia pod rot (Virginia and North Carolina only) <b>FOLIAR:</b> Early leaf spot Late leaf spot Leaf rust Web blotch (Phoma) Pepper spot ( <i>Leptosphaerulina</i> )	7.2 fl oz per acre
	<b>FOUR-APPLICATION SPRAY PROGRAM:</b> Apply the specified rate in a preventive spray schedule. See table below for proper timing of applications. Applications of chlorothalonil should be made prior to and following applications of Quali-Pro Tebuconazole 3.6F to discourage development of resistant strains of fungi. For optimum control of foliar diseases such as leaf rust, web blotch, and pepper spot, the lowest label labeled rate of a spray surfactant should be tank-mixed with Quali-Pro Tebuconazole 3.6F. <b>LEAF SPOT ADVISORY SCHEDULE:</b> For control of soilborne diseases in an advisory schedule, apply Quali-Pro Tebuconazole 3.6F in the first advisory spray in July and continue Quali-Pro Tebuconazole 3.6F applications at 14-day intervals. Applications after August 15 should be tank mixed with chlorothalonil for resistance management purposes.	
<b>GENERAL DIRECTIONS:</b> For optimum control of the specified soilborne diseases, four consecutive applications of Quali-Pro Tebuconazole 3.6F must be made at 14-day intervals. A maximum of 28.8 fluid ounces of Quali-Pro Tebuconazole 3.6F may be applied per crop season. Quali-Pro Tebuconazole 3.6F may be applied up to 14 days before harvest. Do not feed hay or threshings or allow livestock to graze in treated areas. Quali-Pro Tebuconazole 3.6F is a sterol demethylation inhibitor (DMI) fungicide. Chlorothalonil may be tank mixed at the rate of 12 ounces of active ingredient with Quali-Pro Tebuconazole 3.6F as a leaf spot resistance management strategy. A spray surfactant is not necessary when Quali-Pro Tebuconazole 3.6F is tank mixed with chlorothalonil. Mixing or alternating Quali-Pro Tebuconazole 3.6F with other DMI fungicides may lead to resistance. Quali-Pro Tebuconazole 3.6F must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by <i>Sclerotium rolfsii</i> and <i>Rhizoctonia solani</i> . Drought conditions will decrease the effectiveness of Quali-Pro Tebuconazole 3.6F against the root and pod rots. Use Quali-Pro Tebuconazole 3.6F in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, such as proper crop rotation practices.		
<b>Timing of QUALI-PRO TEBUCONAZOLE 3.6F Application for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot</b>		
<b>Spray Program</b>	<b>QUALI-PRO TEBUCONAZOLE 3.6F Application No.</b>	<b>Chlorothalonil Application No.</b>
7 applications	3, 4, 5 and 6	1, 2 and 7
PECAN	Brown leaf spot ( <i>Sirosporium diffusium</i> ) Downy spot ( <i>Mycosphaerella caryigena</i> ) Liver spot ( <i>Gnomonia caryae</i> ) Scab ( <i>Cladosporium caryigenum</i> ) Vein spot ( <i>Gnomonia nerviseda</i> ) Zonate leaf spot ( <i>Grovesinia pyramidalis</i> )	4 to 8 fl oz per acre

	<p><b>Notes:</b> Apply Quali-Pro Tebuconazole 3.6F in a preventive spray schedule beginning at early bud break (young leaves unfolding), and continue applications at 10- to 14-day intervals through the pollination period. Quali-Pro Tebuconazole 3.6F should be applied at 4 fl. oz. per acre in a tank-mix with the labeled rate of Super-Tin® in cover sprays. Follow label directions for the use of Super-Tin. Do not add a surfactant to the spray solution when tank-mixing Quali-Pro Tebuconazole 3.6F with Super-Tin. Apply Quali-Pro Tebuconazole 3.6F in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 to 8 fl. oz. per acre of Quali-Pro Tebuconazole 3.6F to full-size mature trees, and 4 to 6 fl. oz. per acre of Quali-Pro Tebuconazole 3.6F to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases, or when severe disease conditions exist. The lowest labeled rate of a surfactant may be added to the spray solution for optimum control of the indicated diseases. Do not apply after shucks begin to split. A maximum of 32 fl. oz. of Quali-Pro Tebuconazole 3.6F may be applied per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas.</p>	
<p><b>PECAN (continued): General Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Quali-Pro Tebuconazole 3.6F. Quali-Pro Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Quali-Pro Tebuconazole 3.6F will be resistant to weathering. Quali-Pro Tebuconazole 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Restricted-entry interval (REI) = 12 hours.</p>		
<b>SOYBEAN</b>	<p>Rust (<i>Phakopsora pachyrhizi</i>) Powdery mildew (<i>Microsphaera diffusa</i>)</p>	3 to 4 fl oz per acre
<p><b>Use Directions:</b> Apply Quali-Pro Tebuconazole 3.6F as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications on a 10- to 14-day spray interval if environmental conditions are favorable for continued disease development. Use of the higher rates and shorter spray intervals are labeled when disease pressure is severe. The lowest label labeled rate of a spray surfactant must be tank-mixed with Quali-Pro Tebuconazole 3.6F. Quali-Pro Tebuconazole 3.6F should be applied in a minimum for 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment.</p> <p><b>Restrictions:</b> Applications may not be made within 21 days of harvest. Do not apply more than 3 applications per season. Do not apply more than 12 fl. oz/a per use season. Restricted-entry interval (REI) = 12 hours.</p>		
<b>SUNFLOWER</b>	<p>Rust (<i>Puccinia helianthi</i>)</p>	4 to 6 fl oz per acre
<p><b>Notes:</b> Apply specific dosage of Quali-Pro Tebuconazole 3.6F at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Application may be repeated at 14 days if necessary to maintain control of the disease. Apply specified dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Do not apply more than 16 fl. oz. of Quali-Pro Tebuconazole 3.6F F per acre per season or within 50 days of harvest.</p>		
<p><b>General Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Quali-Pro Tebuconazole 3.6F. Contact your state Extension Service or Bayer representative for a list of approved surfactants. Quali-Pro Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Quali-Pro Tebuconazole 3.6F will be resistant to weathering. Quali-Pro Tebuconazole 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.</p>		
<b>TURNIP</b> (Application is limited to East of the Rockies)	<p>Cercospora leaf spot (<i>Cercospora brassicicola</i>)</p>	4 to 7.2 fl. oz. per acre
<p><b>Notes:</b> Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-day intervals. Quali-Pro Tebuconazole 3.6F may be applied up to 7 days before harvest. Do not apply more than 28.8 fl. oz. of Quali-Pro Tebuconazole 3.6F per acre per crop season.</p>		
<p><b>General Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Quali-Pro Tebuconazole 3.6F. Quali-Pro Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Quali-Pro Tebuconazole 3.6 F will be resistant to weathering. Quali-Pro Tebuconazole 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.</p>		
<b>WHEAT</b>	<p>Rusts leaf, stem, and stripe (<i>Puccinia</i> spp.) Head blight or scab (<i>Fusarium</i> spp.) – Suppression</p>	4 fl oz per acre
<p><b>Notes:</b> Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. A maximum of 4 fl. oz. of Quali-Pro Tebuconazole 3.6F may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw may be fed or used for bedding. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with Quali-Pro Tebuconazole 3.6F. Apply Quali-Pro Tebuconazole 3.6F in a minimum of 10 gallons of spray solution per acre by ground, or in a minimum of 5 gallons of spray solution per acre by air.</p> <p><b>Application timing directions:</b> Rusts: Apply Quali-Pro Tebuconazole 3.6 F at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Quali-Pro Tebuconazole 3.6F for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).</p>		
<p><b>General Comments:</b> For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Quali-Pro Tebuconazole 3.6F. Quali-Pro Tebuconazole 3.6F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Quali-Pro Tebuconazole 3.6F will be resistant to weathering. Quali-Pro Tebuconazole 3.6F is a demethylation inhibitor (DMI) fungicide (Group 3).</p>		

<b>SEED TREATMENT- Corn (Sweet Corn, Field Corn, Field Corn Grown For Seed, and Popcorn)</b> For control of soilborne and seedborne Fusarium and soilborne and seedborne head smut.		
<b>SEED LABELING:</b> To meet U.S. Federal Seed Act requirements, all seed treated with Quali-Pro Tebuconazole 3.6F must be labeled:		
<b>TREATED SEED, DO NOT USE FOR FOOD, FEED, OR OIL PURPOSES. Treat with Tebuconazole.</b>		
<b>USE PRECAUTION:</b> When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.		
<b>DISEASE</b>	<b>RATE FL OZ/CWT</b>	<b>DIRECTIONS FOR USE</b>
<b>Soilborne and Seedborne</b> Fusarium	0.071	Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Seed should be sound and well cured prior to treatment. Product should be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates recommended for the crop to be treated with Quali-Pro Tebuconazole 3.6F. The length of control will vary depending on the rate used.
<b>Soilborne and Seedborne</b> Head smut ( <i>Sphacelotheca reiliana</i> )	0.27-0.54	

**OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.**

Apply only during alternate years in fields adjacent to aquatic areas listed above.  
Do not apply by ground or air within 100 feet of aquatic areas listed above.  
Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.  
Spray Drift Management: For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.  
Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.  
Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.  
Make aerial or ground applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.  
Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.  
Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.  
Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

**ROTATIONAL CROPS**

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.  
**PESTICIDE STORAGE:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.  
Handle and open container in a manner as to prevent spillage. If container is leaking invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.  
**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.  
**CONTAINER DISPOSAL:**  
**Nonrefillable Container (five gallons or less):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**LIMITATION OF WARRANTY AND LIABILITY**

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.  
By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES** and **LIMITATIONS OF LIABILITY**.  
**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc. All such risks shall be assumed by the user or buyer.  
**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.  
**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Makhteshim Agan of North America, Inc.'s election, the replacement of product.

Quali-Pro Tebuconazole is a registered trademark of Irvita Plant Protection, N.V.